
# Delirium clinical care standard indicators: 4b-Proportion of patients with delirium who receive a set of interventions to treat the causes of delirium, based on a comprehensive assessment



**Exported from METEOR (AIHW's Metadata Online Registry)** 

#### © Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY4.0 (CC BY4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

## Delirium clinical care standard indicators: 4b-Proportion of patients with delirium who receive a set of interventions to treat the causes of delirium, based on a comprehensive assessment

## Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

**Short name:** Indicator 4b-Proportion of patients with delirium who receive a set of interventions

to treat the causes of delirium, based on a comprehensive assessment

METEOR identifier: 628100

Registration status: Health, Standard 12/09/2016

**Description:** Proportion of patients with <u>delirium</u> who receive a set of interventions to treat the

causes of delirium, based on a comprehensive assessment.

Rationale: A range of risk factors consistently associated with the incidence of delirium have

been identified in the literature (Australian and New Zealand Society for Geriatric Medicine 2012). Some of these may be modifiable, for example, medical illness related factors or laboratory abnormalities such as low albumin and polypharmacy. Identifying underlying causes of delirium can help to target interventions (Ahmed et

al. 2014).

Indicator set: Clinical care standard indicators: delirium

Health, Standard 12/09/2016

Outcome area: <u>Identifying and treating underlying causes</u>

Health, Standard 12/09/2016

## Collection and usage attributes

**Computation description:** For the numerator, the comprehensive assessment of the causes of delirium and

the tailored interventions must be documented in the patient's medical record.

Unless explicitly stated in the indicator, all patients with delirium are included in both the numerator and the denominator, not only those where <u>Hospital service—care</u>

type, code N[N] = 1 Acute care.

Presented as a percentage.

**Computation:** (Numerator ÷ denominator) x 100

**Numerator:** Number of patients with delirium who receive a set of tailored interventions to treat

the causes of delirium, based on a comprehensive assessment.

**Denominator:** Number of patients with delirium.

Comments: This indicator was sourced from The Ontario senior friendly hospital strategy

delirium and functional decline indicators (Wong et al. 2012).

#### Representational attributes

Representation class: Percentage

Data type:RealUnit of measure:PersonFormat:N[NN]

### Source and reference attributes

**Submitting organisation:** Australian Commission on Safety and Quality in Health Care

Reference documents: Ahmed S, Leurent B & Sampson EL 2014. Risk factors for incident delirium among

older people in acute hospital medical units: a systematic review and meta-

analysis. Age and Ageing 43(3):326-33.

Australian and New Zealand Society for Geriatric Medicine 2012. Delirium in older

people: position statement 13.

Wong K, Tsang A, Liu B & Schwartz R 2012. The Ontario senior friendly hospital

strategy delirium and functional decline indicators. Toronto: Ontario Local Health

Integration Network.