

# Severe sepsis (Antimicrobial stewardship clinical care standard)

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Severe sepsis (Antimicrobial stewardship clinical care standard)

## Identifying and definitional attributes

<b>Metadata item type:</b>	Glossary Item
<b>METEOR identifier:</b>	629064
<b>Registration status:</b>	<a href="#">Health</a> , Standard 12/09/2016
<b>Definition:</b>	For the purposes of the Antimicrobial stewardship clinical care standard (ACSQHC 2014), severe sepsis is the systematic response to an infection manifested by organ dysfunction, hypoperfusion or hypotension combined with one or more of the following: fever, tachypnoea, elevated white cell count (Antibiotic Expert Groups 2014).

## Collection and usage attributes

<b>Guide for use:</b>	Severe sepsis may include where <a href="#">Episode of care—principal diagnosis, code (ICD-10-AM 9th edn) ANN{,N[N]}</a> or <a href="#">Episode of care—additional diagnosis, code (ICD-10-AM 9th edn) ANN{,N[N]}</a> include one or more of the following codes:
-----------------------	---

- A02.1 Salmonella sepsis
- A20.7 Sepsis plague
- A21.7 Sepsis tularaemia
- A22.7 Anthrax sepsis
- A23.9 Brucellosis, unspecified (includes Brucella sepsis)
- A24.1 Sepsis melioidosis
- A26.7 Erysipelothrix sepsis
- A28.2 Sepsis extraintestinal yersiniosis
- A32.7 Listerial sepsis
- A39.2 Acute meningococcaemia
- A39.2 to A39.4 Sepsis meningococcal
- A39.3 Chronic meningococcaemia
- A39.4 Meningococcaemia, unspecified
- A40 Streptococcal sepsis
- A40.0 Sepsis due to streptococcus, group A
- A40.1 Sepsis due to streptococcus, group B
- A40.2 Sepsis due to streptococcus, group D
- A40.3 Sepsis due to Streptococcus pneumonia/Pneumococcal sepsis
- A40.8 Other streptococcal sepsis
- A40.9 Streptococcal sepsis, unspecified
- A41 Other sepsis
- A41.0 Sepsis due to Staphylococcus aureus
- A41.1 Sepsis due to other specified staphylococcus / Sepsis due to coagulase-negative staphylococcus
- A41.2 Sepsis due to unspecified staphylococcus
- A41.3 Sepsis due to Haemophilus influenzae
- A41.4 Sepsis due to anaerobes
- A41.50 Sepsis due to unspecified Gram-negative organisms
- A41.52 Sepsis due to Pseudomonas
- A41.58 Sepsis due to other gram neg organisms
- A41.8 Other specified sepsis
- A41.9 Sepsis, unspecified
- A42.7 Sepsis due to actinomycotic
- A48.0 Gas gangrene
- A48.3 Sepsis toxic shock syndrome
- A49.9 Sepsis bacteraemia NOS
- A54.8 Sepsis gonococcal
- B00.7 Sepsis herpesviral
- B37.7 Sepsis candidal
- O03.0 Spontaneous abortion, incomplete, complicated by genital tract and pelvic infection [1]
- O03.5 Spontaneous abortion, complete or unspecified, complicated by genital tract and pelvic infection [1]

- O04.0 Medical abortion, incomplete, complicated by genital tract and pelvic infection [\[1\]](#)
- O04.5 Medical abortion, complete or unspecified, complicated by genital tract and pelvic infection [\[1\]](#)
- O05.0 Other abortion, incomplete, complicated by genital tract and pelvic infection [\[1\]](#)
- O05.5 Other abortion, complete or unspecified, complicated by genital tract and pelvic infection [\[1\]](#)
- O06.0 Unspecified abortion, incomplete, complicated by genital tract and pelvic infection [\[1\]](#)
- O06.5 Unspecified abortion, complete or unspecified, complicated by genital tract and pelvic infection [\[1\]](#)
- O07.0 Failed medical abortion, complicated by genital tract and pelvic infection [\[1\]](#)
- O07.5 Other and unspecified failed attempted abortion, complicated by genital tract and pelvic infection [\[1\]](#)
- O08.0 Genital tract and pelvic infection following abortion and ectopic and molar pregnancy [\[1\]](#)
- O08.2 Embolism following abortion and ectopic and molar pregnancy (septic or septicopyaemic embolism) [\[1\]](#)
- O08.5 Metabolic disorders following abortion and ectopic and molar pregnancy [\[1\]](#)
- O75.3 Sepsis during labour [\[1\]](#)
- O85 Sepsis puerperal [\[1\]](#)
- O88.3 Obstetric pyaemic and septic embolism [\[1\]](#)
- P36.0 Sepsis of newborn due to streptococcus, group B [\[2\]](#)
- P36.1 Sepsis of newborn due to other and unspecified streptococci [\[2\]](#)
- P36.2 Sepsis of newborn due to Staphylococcus aureus [\[2\]](#)
- P36.3 Sepsis of newborn due to other and unspecified staphylococci [\[2\]](#)
- P36.4 Sepsis of newborn due to Escherichia coli [\[2\]](#)
- P36.5 Sepsis of newborn due to anaerobes [\[2\]](#)
- P36.8 Other bacterial sepsis of newborn [\[2\]](#)
- P36.9 Bacterial sepsis of newborn, unspecified [\[2\]](#)
- P37.52 Invasive neonatal candidiasis (includes generalised neonatal candidal sepsis) [\[2\]](#)
- R57.2 Septic shock
- R65.0 [SIRS] of infectious origin without acute organ failure
- R65.1 [SIRS] of infectious origin with acute organ failure (severe sepsis)
- T80.2 Sepsis following infusion, transfusion or therapeutic injection [\[3\]](#)
- T81.42 Sepsis following a procedure [\[3\]](#)
- T88.0 Sepsis following immunisation [\[3\]](#)

[\[1\]](#) Must use additional code starting with 'A' or 'B' in the list above.

[\[2\]](#) Must use additional code R65.1 to identify severe sepsis, or R57.2 to identify septic shock.

[\[3\]](#) Must use additional code starting with 'A', 'B' or 'P' in list above.

## Source and reference attributes

<b>Submitting organisation:</b>	Australian Commission on Safety and Quality in Health Care
<b>Reference documents:</b>	ACSQHC (Australian Commission on Safety and Quality in Health Care) 2014. Antimicrobial stewardship clinical care standard. Sydney: ACSQHC.
	Antibiotic Expert Groups 2014. Therapeutic guidelines: antibiotic. Version 15. Melbourne: Therapeutic Guidelines Limited.

## Relational attributes

<b>Metadata items which use this glossary item:</b>	<a href="#">Antimicrobial stewardship: 1a-Median time to first dose of antibiotics for patients with suspected bacterial meningitis, or for actual or suspected severe sepsis</a> <a href="#">Australian Commission on Safety and Quality in Health Care, Retired</a> 25/01/2022 <a href="#">Health</a> , Standard 12/09/2016
---	--