

Hip fracture (Hip fracture care clinical care standard)

Identifying and definitional attributes

Metadata item type: Glossary Item

METEOR identifier: 629165

Registration status:

- [Health](#), Standard 12/09/2016

Definition: For the purposes of the Hip fracture care clinical care standard (ACSQHC 2015a), hip fracture is a break occurring at the top of the thigh bone (femur), near the pelvis.

Collection and usage attributes

Guide for use: For the purposes of the Indicator specification: hip fracture care clinical care standard (ACSQHC 2015b), hip fracture includes where [Episode of care—principal diagnosis, code \(ICD-10-AM 9th edn\) ANN{.N\[N\]}](#) or [Episode of care—additional diagnosis, code \(ICD-10-AM 9th edn\) ANN{.N\[N\]}](#) is any one of the following:

- S72.00 Fracture of neck of femur, part unspecified
- S72.01 Fracture of intracapsular section of femur
- S72.02 Fracture of upper epiphysis (separation) of femur
- S72.03 Fracture of subcapital section of femur
- S72.04 Fracture of midcervical section of femur (includes transcervical not otherwise specified)
- S72.05 Fracture of base of neck of femur (includes cervicotrochanteric section)
- S72.08 Fracture of other parts of neck of femur (includes fracture of hip not otherwise specified, and head of femur)
- S72.10 Fracture of trochanteric section of femur, unspecified
- S72.11 Fracture of intertrochanteric section of femur
- S72.2 Subtrochanteric fracture.

Only includes episodes where the [Hospital service—care type, code N\[N\]](#) = 1 Acute care.

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: ACSQHC (Australian Commission on Safety and Quality in Health Care) 2015a. Hip fracture care clinical care standard. Sydney: ACSQHC.

ACSQHC 2015b. Indicator specification: hip fracture care clinical care standard. Sydney: ACSQHC.

Relational attributes

Metadata items which use this glossary item: [Hip fracture care clinical care standard indicators: 2a-Evidence of local arrangements for timely and effective pain management for hip fracture](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 2a-Evidence of local arrangements for timely and effective pain management for hip fracture](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 1a-Evidence of local](#)

[arrangements for the management of patients with hip fracture in the emergency department](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 1a-Evidence of local arrangements for the management of patients with hip fracture in the emergency department](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 1b-Proportion of patients with a hip fracture who have had their pre-operative cognitive status assessed](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 1b-Proportion of patients with a hip fracture who have had their pre-operative cognitive status assessed](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 3a-Evidence of orthogeriatric \(or alternative physician or medical practitioner\) management during an admitted patient's hip fracture episode of care](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 3a-Evidence of orthogeriatric \(or alternative physician or medical practitioner\) management during an admitted patient's hip fracture episode of care](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 4a-Proportion of patients with a hip fracture receiving surgery within 48 hours of presentation with the hip fracture](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 4a-Proportion of patients with a hip fracture receiving surgery within 48 hours of presentation with the hip fracture](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5a-Proportion of patients with a hip fracture who are mobilised on day one post hip fracture surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5a-Proportion of patients with a hip fracture who are mobilised on day one post hip fracture surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5b-Proportion of patients with a hip fracture with unrestricted weight bearing status immediately post hip fracture surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5c-Proportion of patients with a hip fracture experiencing a new Stage II or higher pressure injury during their hospital stay](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5c-Proportion of patients with a hip fracture experiencing a new Stage II or higher pressure injury during their hospital stay](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5d-Proportion of patients with a hip fracture returning to pre-fracture mobility](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 5d-Proportion of patients with a hip fracture returning to pre-fracture mobility](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 6a-Proportion of patients with a hip fracture receiving bone protection medicine prior to separation from the hospital at which they underwent surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 6a-Proportion of patients with a hip fracture receiving bone protection medicine prior to separation from the hospital at which they underwent surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 6b-Proportion of patients with a hip fracture readmitted to hospital with another femoral fracture within 12 months of admission from initial hip fracture](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 6b-Proportion of patients with a hip fracture readmitted to hospital with another femoral fracture within 12 months of admission from initial hip fracture](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 7a-Evidence of local arrangements for the development of an individualised care plan for hip fracture patients prior to the patient's separation from hospital](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 7a-Evidence of local arrangements for the development of an individualised care plan for hip fracture patients prior to the patient's separation from hospital](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 7b-Proportion of patients with a hip fracture returning to private residence](#)

- [Australian Commission on Safety and Quality in Health Care](#), Retired 15/05/2018
- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 7b-Proportion of patients with a](#)

[hip fracture returning to private residence](#)

- [Australian Commission on Safety and Quality in Health Care](#), Standard 15/05/2018

[Hip fracture care clinical care standard indicators: 8a-Re-operation of hip fracture patients within 30-day follow-up](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 8a-Re-operation of hip fracture patients within 30-day follow-up](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 8b-Survival at 30 days post-admission for hip fracture surgery](#)

- [Health](#), Standard 12/09/2016

[Hip fracture care clinical care standard indicators: 8b-Survival at 30 days post-admission for hip fracture surgery](#)

- [Health](#), Standard 12/09/2016