

© Australian Institute of Health and Welfare 2024

This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Palliative care phase code N

Identifying and definitional attributes

Metadata item type: Value Domain

METEOR identifier: 639015

Registration status: Independent Hospital Pricing Authority, Standard 03/11/2016

Definition: A code set representing the phases of palliative care.

Representational attributes

Representation class: Code

Data type: Number

Format: N

Maximum character length: 1

ValueMeaningPermissible values:1Stable2Unstable3Deteriorating4TerminalSupplementary values:9Not reported

Collection and usage attributes

Guide for use:

The palliative care phase is the stage of the palliative care patient's illness.

CODE 1 Stable

Patient problems and symptoms are adequately controlled by established plan of care and

- Further interventions to maintain symptom control and quality of life have been planned and
- Family/carer situation is relatively stable and no new issues are apparent.

CODE 2 Unstable

An urgent change in the plan of emergency treatment is required because

- Patient experiences a new problem that was not anticipated in the existing plan of care, and/or
- Patient experiences a rapid increase in the severity of a current problem; and/or
- Family/ carers circumstances change suddenly impacting on patient care.

CODE 3 Deteriorating

The care plan is addressing anticipated needs but requires periodic review because

- · Patients overall functional status is declining and
- · Patient experiences a gradual worsening of existing problemand/or
- Patient experiences a new but anticipated problem and/or
- Family/carers experience gradual worsening distress that impacts on the patient care.

CODE 4 Terminal

Death is likely within days.

CODE 9 Not reported

The phase of the illness has not been reported.

Palliative care phases are not sequential and a patient may move back and forth between phases. Palliative care phases provide a clinical indication of the type of care required and have been shown to correlate strongly with survival within longitudinal prospective studies.

Source and reference attributes

Origin:

Palliative Care Outcomes Collaboration 2014. Palliative Care Outcomes Collaboration Clinical Manual. Australian Health Services Research Institute,

University of Wollongong, Sydney. Viewed 19 May 2016,

http://ahsri.uow.edu.au/content/groups/public/@web/@chsd/@pcoc/

documents/doc/uow129133.pdf

Relational attributes

Related metadata references:

Supersedes Palliative care phase code N

Health, Standard 11/04/2014

Independent Hospital Pricing Authority, Superseded 03/11/2016

Tasmanian Health, Superseded 19/05/2020

Has been superseded by Palliative care phase code N

Aged Care, Standard 30/06/2023 Health, Standard 25/01/2018

Independent Hospital Pricing Authority, Standard 18/01/2018

Tasmanian Health, Standard 19/05/2020

Data elements implementing this value domain:

Episode of admitted patient care—palliative care phase, code N Independent Hospital Pricing Authority, Standard 03/11/2016