

# **Osteoarthritis of the knee clinical care standard indicators: 6a-Proportion of patients with knee osteoarthritis with an agreed date for a review**

**Exported from METEOR (AIHW's Metadata Online Registry)**

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Osteoarthritis of the knee clinical care standard indicators: 6a-Proportion of patients with knee osteoarthritis with an agreed date for a review

## Identifying and definitional attributes

<b>Metadata item type:</b>	Indicator
<b>Indicator type:</b>	Indicator
<b>Short name:</b>	Indicator 6a-Proportion of patients with knee osteoarthritis with an agreed date for a review
<b>METEOR identifier:</b>	644312
<b>Registration status:</b>	<a href="#">Health</a> , Standard 02/08/2017
<b>Description:</b>	Proportion of patients with knee <a href="#">osteoarthritis</a> with an agreed date for a review.
<b>Rationale:</b>	Follow-up should be individualised for each patient, and occur at intervals agreed with the patient (NICE 2014).
<b>Indicator set:</b>	<a href="#">Clinical care standard indicators: osteoarthritis of the knee</a> <a href="#">Health</a> , Standard 02/08/2017

## Collection and usage attributes

<b>Population group age from:</b>	45 years
<b>Computation description:</b>	<p>For the numerator, the agreed date for a review of knee osteoarthritis should be documented (which may be in the form of an appointment made at a date in the future).</p> <p>The review should be within 12 months of initial diagnosis or referral for knee osteoarthritis, and then at agreed intervals thereafter.</p> <p>Presented as a percentage.</p>
<b>Computation:</b>	$(\text{Numerator} \div \text{denominator}) \times 100$
<b>Numerator:</b>	Number of patients with knee osteoarthritis with a documented agreed date for a review.
<b>Denominator:</b>	Number of patients with knee osteoarthritis.
<b>Comments:</b>	This indicator has been adapted from the <i>Osteoarthritis quality standard</i> (NICE 2015).

## Representational attributes

<b>Representation class:</b>	Percentage
<b>Data type:</b>	Real
<b>Unit of measure:</b>	Service event
<b>Format:</b>	N[NN]

## Accountability attributes

<b>Other issues caveats:</b>	Applicable setting: all healthcare settings where care is provided to patients with knee osteoarthritis, including primary care, specialist care, hospitals and community settings.
------------------------------	---

## Source and reference attributes

<b>Submitting organisation:</b>	Australian Commission on Safety and Quality in Health Care
---------------------------------	--

**Reference documents:**

NICE (National Institute for Health and Care Excellence) 2014. Osteoarthritis: The care and management of osteoarthritis in adults. NICE Clinical guideline 177. London: NICE.

NICE 2015. Osteoarthritis Quality standard 87. London: NICE.