

#### © Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

# **Osteoarthritis**

# Identifying and definitional attributes

**Metadata item type:** Glossary Item

METEOR identifier: 644538

**Registration status:** <u>Health</u>, Standard 02/08/2017

**Definition:** A clinical syndrome of joint pain accompanied by varying degrees of functional

limitation and reduced quality of life. Pain, reduced function and effects on a person's ability to carry out their day-to-day activities can be important

consequences. Osteoarthritis is characterised pathologically by localised loss of articular cartilage (cartilage lining the articular ends of the bones), as well as changes to bone and other joint structures and associated inflammation.

### Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

**Reference documents:** NICE (National Institute for Health and Care Excellence) 2014. Osteoarthritis: care

and management in adults. Clinical Guideline 177. London: NICE. Viewed 29

March 2016, https://www.nice.org.uk/guidance/cg177.

## Relational attributes

Metadata items which use this glossary item:

Osteoarthritis of the knee clinical care standard indicators: 1a-Local arrangements to ensure that patients newly diagnosed with knee osteoarthritis have a

comprehensive assessment

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 1b-Proportion of patients newly diagnosed with knee osteoarthritis who have a comprehensive assessment

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 2a-Local arrangements for clinically based diagnosis of knee osteoarthritis without use of imaging for people with knee pain and other symptoms suggestive of osteoarthritis

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 2b-Proportion of patients clinically diagnosed with knee osteoarthritis, without imaging

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 3-Proportion of patients newly diagnosed with knee osteoarthritis who have an individualised self-management plan

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 4a-Proportion of patients newly diagnosed with knee osteoarthritis with a documented recommendation regarding regular exercise

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 4b-Proportion of patients with knee osteoarthritis who were overweight or obese who lost weight

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 5a-Local arrangements to ensure that patients with knee osteoarthritis are prescribed or recommended medicines in accordance with the current Therapeutic guidelines: Rheumatology

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 5b-Proportion of patients with knee osteoarthritis prescribed oral non-steroidal anti-inflammatory drugs (NSAIDs) with documented assessment of risks

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 5c-Proportion of

patients prescribed opioids for longer than three months for the management of pain associated with knee osteoarthritis

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 6a-Proportion of patients with knee osteoarthritis with an agreed date for a review Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 6b-Proportion of patients with knee osteoarthritis with evidence of pain and function assessments within the previous 12 months

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 6c-Proportion of patients with knee osteoarthritis who have documented pain level reduction of at least 20%, 12 months after initiation or change of treatment

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 6d-Proportion of patients with knee osteoarthritis with a functional limitation who have a 10% or greater improvement in function 12 months after initiation or change of treatment Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 7a-Number of patients undergoing arthroscopic procedures for knee osteoarthritis

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 7b-Proportion of patients with knee osteoarthritis referred for consideration of surgery who were supported with non-surgical core treatments for at least three months

Health, Standard 02/08/2017

Osteoarthritis of the knee clinical care standard indicators: 8-Local arrangements to ensure that patients' experience of the care that they receive for knee osteoarthritis is regularly measured and acted upon

Health, Standard 02/08/2017