Alcohol consumption frequency after 20 weeks of pregnancy

Exported from METEOR (AIHW's Metadata Online Registry)
© Australian Institute of Health and Welfare 2024
This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.
You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our

meteor@aihw.gov.au.

attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Alcohol consumption frequency after 20 weeks of pregnancy

Identifying and definitional attributes

Metadata item type: Property
METEOR identifier: 691054

Registration status: <u>Health,</u> Standard 12/12/2018

Tasmanian Health, Standard 09/11/2023

Definition: The frequency of consuming drinks containing alcohol after 20 weeks of pregnancy

until birth.

Context: Perinatal

Property group: <u>Lifestyle characteristics</u>

Source and reference attributes

Submitting organisation: National Perinatal Data Development Committee

Relational attributes

Data Element Concepts

implementing this

property:

Female—alcohol consumption frequency after 20 weeks of pregnancy

Health, Standard 12/12/2018

Tasmanian Health, Standard 09/11/2023