

# Indication for caesarean section code N[N]

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Indication for caesarean section code N[N]

## Identifying and definitional attributes

<b>Metadata item type:</b>	Value Domain
<b>METEOR identifier:</b>	695687
<b>Registration status:</b>	<a href="#">Health</a> , Superseded 03/12/2020
<b>Definition:</b>	A code set representing indications for a caesarean section.

## Representational attributes

<b>Representation class:</b>	Code
<b>Data type:</b>	Number
<b>Format:</b>	N[N]
<b>Maximum character length:</b>	2

	Value	Meaning
<b>Permissible values:</b>	1	Fetal compromise
	2	Suspected fetal macrosomia
	3	Malpresentation
	10	Lack of progress; less than or equal to 3 cm cervical dilatation
	11	Lack of progress in the first stage; greater than 3 cm to less than 10 cm cervical dilatation
	12	Lack of progress in the second stage
	13	Placenta praevia
	14	Placental abruption
	15	Vasa praevia
	16	Antepartum/intrapartum haemorrhage
	17	Multiple pregnancy
	18	Unsuccessful attempt at assisted delivery
	19	Cord prolapse
	20	Previous adverse perinatal outcome
	21	Previous caesarean section
	22	Previous severe perineal trauma
	23	Previous shoulder dystocia
	29	Other obstetric, medical, surgical, psychological indications
	30	Maternal choice in the absence of any obstetric, medical, surgical, psychological indications
<b>Supplementary values:</b>	99	Not stated/inadequately described

## Collection and usage attributes

**Guide for use:****CODE 1 Fetal compromise**

Includes suspected or actual fetal compromise and intra uterine growth restriction.

**CODE 10 Lack of progress; less than or equal to 3 cm cervical dilatation**

Lack of progress includes slow or no progress.

May be appropriate for cases of unsuccessful induction.

**CODE 11 Lack of progress in the first stage; greater than 3 cm to less than 10 cm cervical dilatation**

Lack of progress includes slow or no progress.

**CODE 12 Lack of progress in the second stage**

Lack of progress includes slow or no progress.

**CODE 13 Placenta praevia**

Record placenta praevia as the indication for caesarean section if there is ultrasound or clinical evidence that the edge of the placenta covers the internal cervical os, or encroaches into the lower segment less than 2 cm away from the internal cervical os.

**CODE 14 Placental abruption**

Record placental abruption as the indication for caesarean section if there is ultrasound or clinical evidence antenatally of abruption of the placenta prior to onset or during labour.

**CODE 15 Vasa praevia**

Record vasa praevia as the indication for caesarean section if there is ultrasound or visual evidence of exposed fetal blood vessels running across the fetal membrane below or at the level of the fetal presenting part in the lower segment of the uterus. Recorded when the caesarean section is planned or in the case of an emergency caesarean section when the vessels may have ruptured.

**CODE 16 Antepartum/intrapartum haemorrhage**

Record antepartum/intrapartum haemorrhage as the indication for caesarean section if there has been any antenatal or intrapartum vaginal bleeding that leads to the immediate delivery of the baby by caesarean section. Should only be recorded as a main indication if a more specific cause of the antepartum/intrapartum haemorrhage is not known.

Where there is a vasa praevia and an antepartum/intrapartum haemorrhage, Code 15 is to be recorded as the main indication and Code 16 as an additional indication.

**CODE 22 Previous severe perineal trauma**

Includes previous 4th degree perineal tears and other perineal damage or injury where a female is symptomatic and the clinician determines the caesarean section is justified.

**CODE 29 Other obstetric, medical, surgical, psychological indications**

Where a female has a psychopathological indication for caesarean section, e.g. extreme fear of natural childbirth, this code should be recorded. It is not to be recorded for psychosocial indications which should be recorded as Code 30.

**CODE 30 Maternal choice in the absence of any obstetric, medical, surgical, psychological indications**

Includes psychosocial indications.

Not to be recorded in conjunction with additional indications.

**Comments:** Codes 4 to 9 have been omitted as these codes are no longer in use. For information about their meaning in previous data elements, see superseded versions.

## Source and reference attributes

**Submitting organisation:** National Perinatal Data Development Committee

## Relational attributes

**Related metadata references:** Supersedes [Indication for caesarean section code N\[N\] Health](#), Superseded 12/12/2018  
Has been superseded by [Indication for caesarean section code N\[N\] Health](#), Standard 03/12/2020  
[Tasmanian Health](#), Standard 08/11/2023

**Data elements implementing this value domain:** [Birth event—additional indication for caesarean section, code N\[N\] Health](#), Superseded 03/12/2020  
[Birth event—main indication for caesarean section, code N\[N\] Health](#), Superseded 03/12/2020