

Hip fracture care clinical care standard indicators: 1a- Evidence of local arrangements for the management of patients with hip fracture in the emergency department

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Hip fracture care clinical care standard indicators: 1a-Evidence of local arrangements for the management of patients with hip fracture in the emergency department

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 1a-Evidence of local arrangements for the management of patients with hip fracture in the emergency department
METEOR identifier:	696426
Registration status:	Health , Standard 12/09/2016
Description:	Evidence of local arrangements for the management of patients with hip fracture in the emergency department.
Rationale:	<p>Patients who sustain hip fractures are, and will continue to be, an important patient group in the Australian healthcare system. In most instances, the first hospital contact these patients will have will be when they present to an emergency department (Considine & Hood 2000).</p> <p>Patients presenting to the emergency department with a hip fracture should be rapidly assessed and assigned to a fast track admission pathway (SIGN 2009; Hill et al. 2014). The rate of post-operative complications, including confusion, healthcare associated infection, length of hospital stay and 12 month mortality, is lower in hip fracture patients managed by a comprehensive multidisciplinary fast-track treatment and care programme (SIGN 2009).</p>
Indicator set:	Clinical care standard indicators: hip fracture 2018 Australian Commission on Safety and Quality in Health Care , Standard 15/05/2018
Outcome area:	Care at presentation Health , Standard 12/09/2016

Collection and usage attributes

Computation description:	<p>Documented local arrangements for the management of patients with hip fracture in the emergency department that address timely assessment and management of the patient's medical conditions, including but not limited to:</p> <ul style="list-style-type: none">• diagnostic imaging• pain control• cognitive assessment. <p>The documentation may be in the form of local protocols and/or a clinical pathway.</p>
Computation:	Yes/No

Representational attributes

Representation class:	Count
Data type:	Real
Unit of measure:	Service event
Format:	Yes/No

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Considine J & Hood K 2000. Emergency department management of hip fractures: development of an evidence-based clinical guideline by literature review and consensus. *Emergency Medicine* 12(4):329-36.

Hili S, Dawe EJC, Lindisfame EAO & Stott PM 2014. Perioperative management of elderly patients suffering a hip fracture. *British Journal of Hospital Medicine* 75(2):78-82.

SIGN (Scottish Intercollegiate Guidelines Network) 2009. Management of hip fracture in older people: a national guideline. NHS Quality Improvement Scotland (NHS QIS).