

# Person—weight (measured), total kilograms N[NN].N

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Person—weight (measured), total kilograms N[NN].N

## Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Weight in kilograms (measured)
METEOR identifier:	702085
Registration status:	<a href="#">Health</a> , Standard 12/12/2018 <a href="#">Tasmanian Health</a> , Standard 24/03/2023
Definition:	The weight (body mass) of a person measured in kilograms.
Data Element Concept:	<a href="#">Person—weight</a>
Value Domain:	<a href="#">Total kilograms N[NN].N</a>

## Value domain attributes

## Representational attributes

Representation class:	Total	
Data type:	Number	
Format:	N[NN].N	
Maximum character length:	4	
	Value	Meaning
Supplementary values:	999.9	Not collected
Unit of measure:	Kilogram (Kg)	

## Collection and usage attributes

Guide for use:	A continuous variable measured to the nearest 0.1 kg.
	CODE 999.9 Not collected
	Use this code if measured weight is not collected.

## Data element attributes

## Collection and usage attributes

Guide for use:	In order to ensure consistency in measurement, the measurement protocol described under Collection methods should be used.
Collection methods:	<p>The collection of anthropometric measurements, particularly in those who are overweight or obese or who are concerned about their weight, should be performed with great sensitivity and without drawing attention to an individual's weight.</p> <p>The measurement protocol described below is that recommended by the WHO Expert Committee (1995).</p> <p>Measurement protocol:</p> <p>Equipment used should be described and reported. Scales should have a resolution of at least 0.1kg and should have the capacity to weigh up to at least 200 kg. Measurement intervals and labels should be clearly readable under all conditions of use of the instrument. Scales should be capable of being calibrated across the entire range of measurements. Precision error should be no more than</p>

0.1 kg. Scales should be calibrated on each day of use. Manufacturers' guidelines should be followed with regard to the transportation of the scales.

Adults and children who can stand:

The subject stands over the centre of the weighing instrument, with the body weight evenly distributed between both feet.

Heavy jewellery should be removed and pockets emptied. Light indoor clothing can be worn, excluding shoes, belts, and sweater. Any variations from light indoor clothing (e.g. heavy clothing, such as kaftans or coats worn because of cultural practices) should be noted on the data collection form. Adjustments for non-standard clothing (i.e. other than light indoor clothing) should only be made in the data checking/cleaning stage prior to data analysis.

If the subject has had one or more limbs amputated, record this on the data collection form and weigh them as they are. If they are wearing an artificial limb, record this on the data collection form but do not ask them to remove it. Similarly, if they are not wearing the limb, record this but do not ask them to put it on.

The measurement is recorded to the nearest 0.1 kg. If the scales do not have a digital readout, take a repeat measurement. If the two measurements disagree by more than 0.5 kg, then take a third measurement. All raw measurements should be recorded on the data collection form. If practical, it is preferable to enter the raw data into the database as this enables intra-observer and, where relevant, inter-observer errors to be assessed. The subject's measured weight is subsequently calculated as the mean of the two observations, or the mean of the two closest measurements if a third is taken, and recorded on the form. If only a mean value is entered into the database then the data collection forms should be retained.

It may be necessary to round the mean value to the nearest 0.1 kg. If so, rounding should be to the nearest even digit to reduce systematic over reporting (Armitage and Berry 1994). For example, a mean value of 72.25 kg would be rounded to 72.2 kg, while a mean value of 72.35 kg would be rounded to 72.4 kg.

Infants:

Birth weight and gender should be recorded with gestational age. During infancy a levelled pan scale with a bean and movable weights or digital scales capable of measuring to two decimal places of a kilogram are acceptable. Birth weight should be determined within 12 hours of birth. The infant, with or without a nappy or diaper is placed on the scales so that the weight is distributed equally about the centre of the pan. When the infant is lying or suspended quietly, weight is recorded to the nearest 10 grams. If the nappy or diaper is worn, its weight is subtracted from the observed weight i.e. reference data for infants are based on nude weights.

Validation and quality control measures:

If practical, equipment should be checked daily using one or more objects of known weight in the range to be measured. It is recommended that the scale be calibrated at the extremes and in the mid range of the expected weight of the population being studied.

Within- and, if relevant, between-observer variability should be reported. They can be assessed by the same (within-) or different (between-) observers repeating the measurement of weight, on the same subjects, under standard conditions after a short time interval. The standard deviation of replicate measurements (technical error of measurement) between observers should not exceed 0.5 kg and be less than 0.5 kg within observers.

Extreme values at the lower and upper end of the distribution of measured height should be checked both during data collection and after data entry. Individuals should not be excluded on the basis of true biological difference.

Last digit preference, and preference or avoidance of certain values, should be analysed in the total sample and (if relevant) by observer, survey site and over time if the survey period is long.

<b>Comments:</b>	<p>This metadata item applies to persons of all ages. It is recommended for use in population surveys and health care settings.</p> <p>It is recommended that in population surveys, sociodemographic data including ethnicity should be collected, as well as other risk factors including physiological status (e.g. pregnancy), physical activity, smoking and alcohol consumption. Summary statistics may need to be adjusted for these variables.</p> <p>Metadata items currently exist for sex, date of birth, country of birth, Indigenous status and smoking. Metadata items are being developed for physical activity.</p> <p>Presentation of data:</p> <p>Means and 95% confidence intervals, medians and centiles should be reported to one decimal place. Where the sample permits, population estimates should be presented by sex and 5-year age groups. However 5-year age groups are not generally suitable for children and adolescents. Estimates based on sample surveys may need to take into account sampling weights.</p> <p>For consistency with conventional practice, and for current comparability with international data sets, recommended centiles are 5, 10, 15, 25, 50, 75, 85, 90 and 95. To estimate the 5th and 95th centiles, a sample size of at least 200 is recommended for each group for which the centiles are being specified.</p> <p>For some reporting purposes, it may be desirable to present weight data in categories. It is recommended that 5 kg groupings are used for this purpose. Weight data should not be rounded before categorisation. The following categories may be appropriate for describing the weights of Australian men, women, children and adolescents, although the range will depend on the population.</p> <p>Weight &lt; 10 kg</p> <p>10 kg = Weight &lt;15 kg</p> <p>15 kg = Weight &lt; 20 kg</p> <p>... in 5 kg categories</p> <p>135 kg = Weight &lt; 140 kg</p> <p>Weight =&gt; 140 kg</p>
------------------	---

## Source and reference attributes

<b>Submitting organisation:</b>	World Health Organization The consortium to develop standard methods for the collection and collation of anthropometric data in children as part of the National Food and Nutrition Monitoring and Surveillance Project, funded by the Commonwealth Department of Health and Ageing
<b>Reference documents:</b>	<p>Clinical Guidelines on the Identification, Evaluation and Treatment of Overweight and Obesity in Adults (US National Heart, Lung and Blood Institute (NHLBI) in cooperation with the National Institute of Diabetes and Digestive and Kidney Diseases).</p> <p>Chronic Diseases and Associated Risk Factors in Australia 2001 (AIHW).</p>

## Relational attributes

<b>Related metadata references:</b>	Supersedes <a href="#">Person—weight (measured), total kilograms N[NN].N Health</a> , Superseded 12/12/2018
<b>Implementation in Data Set Specifications:</b>	<p><a href="#">Perinatal NBEDS 2019–20 Health</a>, Superseded 20/11/2019</p> <p><b>Implementation start date:</b> 01/07/2019</p> <p><b>Implementation end date:</b> 30/06/2020</p> <p><b>DSS specific information:</b></p>

Weight rounded to the nearest whole kilogram is acceptable.

Weight should be recorded in the first trimester, preferably as a measured weight.

However, the female's self-reported weight around the time of conception is acceptable if a measured weight is not available.

It is preferable to collect and record a pregnant female's weight as a measured weight. Where a measured weight has not been provided or it is not possible to ascertain whether the weight has been measured or self-reported, the value 999.9 should be recorded against this data element. Data should then be recorded against the [Person—weight \(self-reported\), total kilograms NN\[N\]](#) data element.

#### [Perinatal NBEDS 2020–21](#)

[Health](#), Superseded 03/12/2020

**Implementation start date:** 01/07/2020

**Implementation end date:** 30/06/2021

**DSS specific information:**

Weight rounded to the nearest whole kilogram is acceptable.

Weight should be recorded in the first trimester, preferably as a measured weight. However, the female's self-reported weight around the time of conception is acceptable if a measured weight is not available.

It is preferable to collect and record a pregnant female's weight as a measured weight. Where a measured weight has not been provided or it is not possible to ascertain whether the weight has been measured or self-reported, the value 999.9 should be recorded against this data element. Data should then be recorded against the [Person—weight \(self-reported\), total kilograms NN\[N\]](#) data element.

#### [Perinatal NBEDS 2021–22](#)

[Health](#), Superseded 17/12/2021

**Implementation start date:** 01/07/2021

**Implementation end date:** 30/06/2022

**DSS specific information:**

This data element is recorded for the mother only.

Weight should be recorded in the first trimester, preferably as a measured weight. However, the female's self-reported weight around the time of conception is acceptable if a measured weight is not available.

It is preferable to collect and record a pregnant female's weight as a measured weight. Where a measured weight has not been provided or it is not possible to ascertain whether the weight has been measured or self-reported, the value 999.9 should be recorded against this data element. Data should then be recorded against the [Person—weight \(self-reported\), total kilograms NN\[N\]](#) data element.

Weight rounded to the nearest whole kilogram is acceptable.

#### [Perinatal NBEDS 2022–23](#)

[Health](#), Superseded 09/12/2022

**Implementation start date:** 01/07/2022

**Implementation end date:** 30/06/2023

**DSS specific information:**

This data element is recorded for the mother only.

Weight should be recorded in the first trimester, preferably as a measured weight. However, the female's self-reported weight around the time of conception is acceptable if a measured weight is not available.

It is preferable to collect and record a pregnant female's weight as a measured weight. Where a measured weight has not been provided or it is not possible to ascertain whether the weight has been measured or self-reported, the value 999.9 should be recorded against this data element. Data should then be recorded against the [Person—weight \(self-reported\), total kilograms NN\[N\]](#) data element.

Weight rounded to the nearest whole kilogram is acceptable.

#### [Perinatal NBEDS 2023–24](#)

[Health](#), Superseded 06/12/2023

**Implementation start date:** 01/07/2023

**Implementation end date:** 30/06/2024

**DSS specific information:**

This data element is recorded for the mother only.

Weight should be recorded in the first trimester, preferably as a measured weight. However, the female's self-reported weight around the time of conception is acceptable if a measured weight is not available.

It is preferable to collect and record a pregnant female's weight as a measured weight. Where a measured weight has not been provided or it is not possible to ascertain whether the weight has been measured or self-reported, the value 999.9 should be recorded against this data element. Data should then be recorded against the [Person—weight \(self-reported\), total kilograms NN\[N\]](#) data element.

Weight rounded to the nearest whole kilogram is acceptable.

[Perinatal NBEDS 2024–25](#)

[Health](#), Standard 06/12/2023

**Implementation start date:** 01/07/2024

**Implementation end date:** 30/06/2025

**DSS specific information:**

This data element is recorded for the mother only.

Weight should be recorded in the first trimester, preferably as a measured weight. However, the female's self-reported weight around the time of conception is acceptable if a measured weight is not available.

It is preferable to collect and record a pregnant female's weight as a measured weight. Where a measured weight has not been provided or it is not possible to ascertain whether the weight has been measured or self-reported, the value 999.9 should be recorded against this data element. Data should then be recorded against the [Person—weight \(self-reported\), total kilograms NN\[N\]](#) data element.

Weight rounded to the nearest whole kilogram is acceptable.