

# Primary cause of perinatal death

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Primary cause of perinatal death

## Identifying and definitional attributes

<b>Metadata item type:</b>	Property
<b>METEOR identifier:</b>	716423
<b>Registration status:</b>	<a href="#">Health</a> , Standard 01/02/2023
<b>Definition:</b>	The single most important factor which led to the chain of events which resulted in a perinatal death.

## Source and reference attributes

**Submitting organisation:** Australian Institute of Health and Welfare

## Relational attributes

**Related metadata references:** See also [Primary cause of neonatal death](#)  
[Health](#), Standard 07/02/2023

**Data Element Concepts implementing this property:** [Product of birth—primary cause of perinatal death](#)  
[Health](#), Standard 07/02/2023