

## © Australian Institute of Health and Welfare 2024

This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

## Primary cause of neonatal death

## Identifying and definitional attributes

Metadata item type: Property **METEOR** identifier: 716427

Registration status: Health, Standard 07/02/2023

**Definition:** The single most important factor in the **neonatal** period which caused a death.

## Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare

**Relational attributes** 

Related metadata See also Primary cause of perinatal death

Health, Standard 01/02/2023 references:

**Data Element Concepts** 

implementing this

Product of birth—primary cause of neonatal death

Health, Standard 07/02/2023 property: