

Acute stroke clinical care standard indicators: 4b- Proportion of patients with a final diagnosis of acute stroke assessed for ongoing rehabilitation using a structured assessment tool prior to separation from acute care, 2019-

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute stroke clinical care standard indicators: 4b- Proportion of patients with a final diagnosis of acute stroke assessed for ongoing rehabilitation using a structured assessment tool prior to separation from acute care, 2019-

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 4b- Proportion of patients with a final diagnosis of acute stroke assessed for ongoing rehabilitation using a structured assessment tool prior to separation from acute care
METEOR identifier:	719058
Registration status:	Australian Commission on Safety and Quality in Health Care , Qualified 09/09/2019
Description:	Proportion of patients with a final diagnosis of acute stroke who are assessed for ongoing stroke rehabilitation using a structured assessment tool prior to separation from acute care.
Indicator set:	Clinical care standard indicators: acute stroke Australian Commission on Safety and Quality in Health Care , Standard 03/11/2020

Collection and usage attributes

Computation description:	<p>A structured assessment tool such as the Australian Stroke Coalition Rehabilitation Working Group. Assessment for Rehabilitation: Pathway and Decision-Making Tool should be used and the outcome of the assessment must be documented in the medical record.</p> <p>Both the numerator and the denominator include patients with a final diagnosis of acute stroke. The final diagnosis is made at the hospital where the patient is admitted for the acute phase of management of their stroke.</p> <p>Separation from acute care is specified as either being transferred to another ward or discharge to the community or home including an aged care facility.</p>
Computation:	$(\text{Numerator} \div \text{denominator}) \times 100$
Numerator:	Number of patients with a final diagnosis of acute stroke who were assessed for rehabilitation using a structured assessment tool prior to discharge from the acute care
Denominator:	Number of patients with a final diagnosis of acute stroke

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Episode
Format:	N[NN]

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
Reference documents:	Australian Stroke Coalition Rehabilitation Working Group 2012. Assessment for Rehabilitation: Pathway and decision-making tool. Melbourne.

