

Acute coronary syndromes clinical care standard indicators: 6a-Proportion of patients with acute coronary syndrome (ACS) provided with an individualised care plan, 2019-

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute coronary syndromes clinical care standard indicators: 6a-Proportion of patients with acute coronary syndrome (ACS) provided with an individualised care plan, 2019-

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 6a-Proportion of ACS patients provided with a written, individualised care plan
METEOR identifier:	719398
Registration status:	Australian Commission on Safety and Quality in Health Care, Qualified 17/09/2019
Description:	Proportion of acute coronary syndrome (ACS) patients provided with a written, individualised care plan (addressing factors such as gradual physical activity, smoking cessation and therapies addressing psychosocial needs).
Rationale:	
Indicator set:	Clinical care standard indicators: acute coronary syndromes Australian Commission on Safety and Quality in Health Care, Standard 29/10/2020

Collection and usage attributes

Computation description:	<p>Both the numerator and the denominator include patients with acute chest pain, or other symptoms suggestive of ACS.</p> <p>Both the numerator and the denominator exclude patients who died in hospital, patients separated on a palliative care pathway, or where adherence to a secondary prevention plan is not indicated.</p> <p>For an individualised care plan to be included for a patient in the numerator, it needs to be a written plan, addressing factors specific to the patient, such as:</p> <ul style="list-style-type: none">• lifestyle counselling on smoking cessation, ambulatory exercise and healthy eating• an angina symptom management plan which entails guidance on the use of prescribed medicines to manage acute worsening or new unstable angina symptoms, and when and how (i.e. call ambulance rather than self-drive) to present to EDs for acute care. <p>Presented as a percentage.</p>
Computation:	$(\text{Numerator} \div \text{denominator}) \times 100$
Numerator:	Number of ACS patients provided with an individualised care plan.
Denominator:	Number of ACS patients separated from hospital.
Comments:	Ensure the advice provided in the care plan is culturally and linguistically appropriate with particular consideration among Aboriginal and Torres Strait Islander patients and CALD groups.

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Episode

Format:

N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: National Heart Foundation of Australia & Cardiac Society of Australia and New Zealand 2016 Executive Working Group 2016. Australian clinical guidelines for the management of acute coronary syndromes 2016. Heart, Lung, and Circulation 25: 895-951.