

# **Acute coronary syndromes clinical care standard indicators: 6b-Proportion patients with acute coronary syndrome (ACS) on aspirin or dual antiplatelet therapy on separation from hospital, 2019-**

**Exported from METEOR (AIHW's Metadata Online Registry)**

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Acute coronary syndromes clinical care standard indicators: 6b-Proportion patients with acute coronary syndrome (ACS) on aspirin or dual antiplatelet therapy on separation from hospital, 2019-

## Identifying and definitional attributes

<b>Metadata item type:</b>	Indicator
<b>Indicator type:</b>	Indicator
<b>Short name:</b>	Indicator 6b-Proportion of patients with ACS discharged on aspirin or dual antiplatelet therapy
<b>METEOR identifier:</b>	719400
<b>Registration status:</b>	<a href="#">Australian Commission on Safety and Quality in Health Care</a> , Qualified 17/09/2019
<b>Description:</b>	Proportion of patients with a final diagnosis of <a href="#">acute coronary syndrome (ACS)</a> who are prescribed aspirin or dual antiplatelet therapy on separation from hospital.
<b>Indicator set:</b>	<a href="#">Clinical care standard indicators: acute coronary syndromes</a> <a href="#">Australian Commission on Safety and Quality in Health Care</a> , Standard 29/10/2020

## Collection and usage attributes

<b>Computation description:</b>	Both the numerator and the denominator include patients with acute chest pain, or other symptoms suggestive of ACS.  The denominator excludes ACS patients who died in hospital, who are being treated with anticoagulants, and for whom aspirin and other antiplatelet therapies are contraindicated, and for whom the contraindication is documented. Contraindications may include advance care directives, being on a palliative care pathway, and clinical judgement, subject to discussion with patients, family and carers.  Presented as a percentage.
<b>Computation:</b>	$(\text{Numerator} \div \text{denominator}) \times 100$
<b>Numerator:</b>	Number of patients with a final diagnosis of ACS who are prescribed aspirin or other dual antiplatelet therapy on separation from hospital.
<b>Denominator:</b>	Number of patients with a final diagnosis of ACS who are separated from hospital.

## Representational attributes

<b>Representation class:</b>	Percentage
<b>Data type:</b>	Real
<b>Unit of measure:</b>	Episode
<b>Format:</b>	N[NN]

## Source and reference attributes

<b>Submitting organisation:</b>	Australian Commission on Safety and Quality in Health Care
<b>Reference documents:</b>	National Heart Foundation of Australia & Cardiac Society of Australia and New Zealand 2016 Executive Working Group 2016. Australian clinical guidelines for the management of acute coronary syndromes 2016. Heart, Lung, and Circulation 25: 895-951.

