

Acute coronary syndromes clinical care standard indicators: 6e-Proportion of patients with reduced left ventricular ejection fraction(LVEF) prescribed a beta blocker and angiotensin-converting enzyme (ACE) inhibitor on separation from hospital, 2019-

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute coronary syndromes clinical care standard indicators: 6e-Proportion of patients with reduced left ventricular ejection fraction(LVEF) prescribed a beta blocker and angiotensin-converting enzyme (ACE) inhibitor on separation from hospital, 2019-

Identifying and definitional attributes

| | |
|-----------------------------|--|
| Metadata item type: | Indicator |
| Indicator type: | Indicator |
| Short name: | Indicator 6e-Proportion of patients with reduced LVEF prescribed a beta blocker and ACE inhibitor or angiotensin receptor blocker (ARB) |
| METEOR identifier: | 719406 |
| Registration status: | Australian Commission on Safety and Quality in Health Care , Qualified 17/09/2019 |
| Description: | Proportion of patients with a final diagnosis of myocardial infarction (MI) and left ventricular ejection fraction less than 40% prescribed a beta blocker and ACE inhibitor or ARB on separation from hospital. |
| Indicator set: | Clinical care standard indicators: acute coronary syndromes Australian Commission on Safety and Quality in Health Care , Standard 29/10/2020 |

Collection and usage attributes

| | |
|---------------------------------|--|
| Computation description: | Both the numerator and the denominator include patients with MI and left ventricular ejection fraction less than 40%. The denominator excludes patients who died in hospital, and for whom beta blockers, ACE inhibitors or ARB are contraindicated and the contraindication is documented. Presented as a percentage. |
| Computation: | $(\text{Numerator} \div \text{denominator}) \times 100$ |
| Numerator: | Number of patients with a final diagnosis of MI and LVEF less than 40% prescribed a beta blocker and ACE inhibitor(or ARB) on separation from hospital. |
| Denominator: | Number of patients with a final diagnosis of MI and LVEF less than 40%. |

Representational attributes

| | |
|------------------------------|------------|
| Representation class: | Percentage |
| Data type: | Real |
| Unit of measure: | Episode |
| Format: | N[NN] |

Source and reference attributes

| | |
|---------------------------------|--|
| Submitting organisation: | Australian Commission on Safety and Quality in Health Care |
|---------------------------------|--|