Acute coronary syndromes clinical care standard indicators: 6f-Proportion of patients with acute coronary syndrome referred to cardiac rehabilitation or other secondary prevention program, 2019-



© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute coronary syndromes clinical care standard indicators: 6f-Proportion of patients with acute coronary syndrome referred to cardiac rehabilitation or other secondary prevention program, 2019-

Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

Short name: Indicator 6f-Proportion of patients referred to cardiac rehabilitation or other

secondary prevention program

METEOR identifier: 719409

Registration status: Australian Commission on Safety and Quality in Health Care, Qualified 17/09/2019

Description: Proportion of patients with an <u>acute coronary syndrome (ACS)</u> with a

documented referral to a cardiac rehabilitation or an alternative secondary

prevention program prior to seperation from hospital.

Indicator set: Clinical care standard indicators: acute coronary syndromes

Australian Commission on Safety and Quality in Health Care, Standard

29/10/2020

Collection and usage attributes

Computation description: Both the numerator and the denominator include patients with acute chest pain, or

other symptoms suggestive of ACS.

Both the numerator and denominator exclude patients who died in hospital, patients for whom cardiac rehabilitation or other secondary prevention program are contraindicated (where the contraindication is documented in the patient's medical record). Contraindications may include advance care directives, being on a palliative care pathway, and clinical judgement, subject to discussion with patients,

family and carers.

Presented as a percentage.

Computation: (Numerator ÷ denominator) x 100

Numerator: Number of patients with a final diagnosis of ACS with a documented referral to a

cardiac rehabilitation or an alternative secondary prevention program.

Denominator: Number of patients with a final diagnosis of ACS separated from hospital.

Comments: The World Health Organization (1993) has defined cardiac rehabilitation as:

"the sum of activities required to influence favourably the underlying cause of the disease, as well as the best possible, physical, mental and social conditions, so that they (people) may, by their own efforts preserve or resume when lost, as normal a place as possible in the community. Rehabilitation cannot be regarded as an isolated form or stage of therapy but must be integrated within secondary

prevention services of which it forms only one face."

For additional information on cardiac rehabilitation, see the *Recommended* Framework for Cardiac Rehabilitation (National Heart Foundation of Australia and

Australian Cardiac Rehabilitation Association 2004).

Representational attributes

Representation class: Percentage

Data type: Real

Unit of measure: Episode Format: N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: National Heart Foundation of Australia and Australian Cardiac Rehabilitation

Association 2004. Recommended framework for cardiac rehabilitation '04.

Canberra: NHFA and ACRA.

WHO (World Health Organization) 1993. Needs and action priorities in cardiac rehabilitation and secondary prevention in patients with CHD. Geneva: WHO.