

Acute coronary syndromes clinical care standard indicators: 6g- Proportion of patients with acute coronary syndrome whose discharge summary was provided to their general practitioner or ongoing clinical provider within 48 hours of separation from hospital

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 6g- Proportion of patients whose discharge summary is provided to their general practitioner or ongoing clinical provider within 48 hours of separation from hospital
METEOR identifier:	719411
Registration status:	<ul style="list-style-type: none">• Australian Commission on Safety and Quality in Health Care, Qualified 17/09/2019

Description: Proportion of patients with a final diagnosis of [acute coronary syndrome \(ACS\)](#) whose discharge summary was provided to their general practitioner (GP) or ongoing clinical provider within 48 hours of separation from hospital.

Indicator set: [Clinical care standard indicators: acute coronary syndromes](#)
[Australian Commission on Safety and Quality in Health Care](#), Standard 29/10/2020

Collection and usage attributes

Computation description: Both the numerator and the denominator include patients with acute chest pain, or other symptoms suggestive of ACS.

Presented as a percentage.

Computation: $(\text{Numerator} \div \text{denominator}) \times 100$

Numerator: Number of patients with a final diagnosis of ACS whose discharge summary was provided to their GP or relevant health service provider within 48 hours of separation from hospital.

Denominator: Number of patients with a final diagnosis of ACS separated from hospital.

Representational attributes

Representation class: Percentage

Data type: Real

Unit of measure: Episode

Format: N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care