Acute coronary syndromes clinical care standard indicators: 6g-Proportion of patients with acute coronary syndrome whose discharge summary was provided to their general practitioner or ongoing clinical provider within 48 hours of separation from hospital



© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute coronary syndromes clinical care standard indicators: 6g-Proportion of patients with acute coronary syndrome whose discharge summary was provided to their general practitioner or ongoing clinical provider within 48 hours of separation from hospital

Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

Short name: Indicator 6g- Proportion of patients whose discharge summary is provided to their

general practitioner or ongoing clinical provider within 48 hours of separation from

hospital

METEOR identifier: 719411

Registration status: Australian Commission on Safety and Quality in Health Care, Qualified 17/09/2019

Description: Proportion of patients with a final diagnosis of <u>acute coronary</u>

syndrome (ACS) whose discharge summary was provided to their general practitioner (GP) or ongoing clinical provider within 48 hours of separation from

hospital.

Indicator set: Clinical care standard indicators: acute coronary syndromes

Australian Commission on Safety and Quality in Health Care, Standard

29/10/2020

Collection and usage attributes

Computation description: Both the numerator and the denominator include patients with acute chest pain, or

other symptoms suggestive of ACS.

Presented as a percentage.

Computation: (Numerator ÷ denominator) x 100

Numerator: Number of patients with a final diagnosis of ACS whose discharge summary

was provided to their GP or relevant health service provider within 48 hours of

separation from hospital.

Denominator: Number of patients with a final diagnosis of ACS separated from hospital.

Representational attributes

Representation class: Percentage

Data type: Real
Unit of measure: Episode
Format: N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care