# Third and Fourth Degree Perineal Tears Clinical Care Standard: 2-Proportion of women who had a vaginal birth who received warm compresses during the second stage of labour



### © Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

## Third and Fourth Degree Perineal Tears Clinical Care Standard: 2-Proportion of women who had a vaginal birth who received warm compresses during the second stage of labour

### Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

**Short name:** 2-Proportion of women who had a vaginal birth who received warm compresses

during the second stage of labour

METEOR identifier: 728357

Registration status: Australian Commission on Safety and Quality in Health Care, Standard 20/04/2021

**Description:** Proportion of women who had a vaginal birth who received warm compresses

during the second stage of labour.

Indicator set: Clinical care standard indicators: Third and Fourth Degree Perineal Tears

Australian Commission on Safety and Quality in Health Care, Standard

20/04/2021

### Collection and usage attributes

Population group age

from:

18 yrs or older

Computation description: For the numerator, 'received' means that it is documented in the medical record

that the woman received warm compresses during the second stage of labour.

Presented as a percentage.

**Computation:** (Numerator ÷ Denominator) x 100

**Numerator:** The number of women who had a vaginal birth who received warm compresses

during the second stage of labour.

**Denominator:** The number of women who had a vaginal birth.

Include separations where an outcome of delivery was record using the ICD codes in Table A, and a caesarean delivery was not recorded (Table B).

Table A: ICD-10-AM codes for delivery outcome

Code	Description
Z37.0	Single live birth
Z37.1	Single stillbirth
Z37.2	Twins, both liveborn
Z37.3	Twins, one liveborn and one stillborn
Z37.4	Twins, both stillborn
Z37.5	Other multiple births, all liveborn
Z37.6	Other multiple births, some liveborn
Z37.7	Other multiple births, all stillborn
Z37.9	Outcome of delivery, unspecified

Table B: ACHI codes for caesarean section

Code	Description
16520-00	Elective classical caesarean section
16520-01	Emergency classical caesarean section
16520-02	Elective lower segment caesarean section
16520-03	Emergency lower segment caesarean section
16520-04	Elective caesarean section, not elsewhere classified
16520-05	Emergency caesarean section, not elsewhere classified

## Representational attributes

Representation class: Percentage

Data type:RealUnit of measure:PersonFormat:N[NN]

### Source and reference attributes

**Submitting organisation:** Australian Commission on Safety and Quality in Health Care

Reference documents: Australian Commission on Safety and Quality in Health Care 2017. National Safety

and Quality Health Service Standards. 2nd ed. Australian Commission on Safety

and Quality in Health Care, Sydney.

Australian Commission on Safety and Quality in Health Care 2021. Third and Fourth Degree Perineal Tears clinical care standard. Australian Commission on

Safety and Quality in Health Care, Sydney.