

Delirium clinical care standard indicators: 5b- Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Delirium clinical care standard indicators: 5b- Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	5b-Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium
METEOR identifier:	745820
Registration status:	Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021
Description:	Proportion of patients with delirium who received multicomponent interventions to treat and manage delirium.
Indicator set:	Clinical care standard indicators: delirium 2021 Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021

Collection and usage attributes

Computation description:	For the numerator and denominator identify patients with delirium using the ICD codes specified in the Hospital-Acquired Complications (HACs) List - Specifications , worksheet 1, where complication group = 11. <i>Delirium</i> . For the numerator, the multicomponent interventions to treat and manage delirium must be documented in the patient's medical record and include tailored interventions to treat any identified cause/s of delirium. Presented as a percentage.
Computation:	$(\text{Numerator} \div \text{Denominator}) \times 100$
Numerator:	Number of patients with delirium who received multicomponent interventions to treat and manage delirium.
Denominator:	The number of patients with delirium.
Comments:	This indicator is based on the Ontario Senior Friendly Hospital Strategy Delirium and Functional Decline Indicators (Wong et al, 2012).

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Person
Format:	N[NN]

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
Reference documents:	Australian Commission on Safety and Quality in Health Care. Delirium Clinical Care Standard. Sydney: ACSQHC; 2021. Wong K, Tsang A, Liu B & Schwartz R. The Ontario Senior Friendly Hospital Strategy Delirium and Functional Decline Indicators. Toronto: Ontario Local Health Integration Network; 2012.

