

Delirium clinical care standard indicators: 6a- Proportion of patients with delirium who were assessed for risk of functional decline, dehydration, malnutrition, falls, and pressure injuries

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Delirium clinical care standard indicators: 6a-Proportion of patients with delirium who were assessed for risk of functional decline, dehydration, malnutrition, falls, and pressure injuries

Identifying and definitional attributes

| | |
|-----------------------------|---|
| Metadata item type: | Indicator |
| Indicator type: | Indicator |
| Short name: | 6a-Proportion of patients with delirium who were assessed for risk of functional decline, dehydration, malnutrition, falls, and pressure injuries |
| METEOR identifier: | 745822 |
| Registration status: | Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021 |
| Description: | Proportion of patients with delirium who were assessed for risk of functional decline, dehydration, malnutrition, falls, and pressure injuries. |
| Indicator set: | Clinical care standard indicators: delirium 2021 Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021 |

Collection and usage attributes

| | |
|---------------------------------|---|
| Computation description: | For the numerator and denominator identify patients with delirium using the ICD codes specified in the Hospital-Acquired Complications (HACs) List - Specifications , worksheet 1, where complication group = 11. <i>Delirium</i> . The numerator includes patients with delirium who were assessed for risk of functional decline, dehydration, malnutrition, falls, and pressure injuries. Presented as a percentage. |
| Computation: | $(\text{Numerator} \div \text{Denominator}) \times 100$ |
| Numerator: | The number of patients with delirium who were assessed for risk of functional decline, dehydration, malnutrition, falls, and pressure injuries |
| Denominator: | The number of patients with delirium. |
| Comments: | This indicator is based on the Key Principles for Care of Confused Hospitalised Older Persons (ACI, 2014). |

Representational attributes

| | |
|------------------------------|------------|
| Representation class: | Percentage |
| Data type: | Real |
| Unit of measure: | Person |
| Format: | N[NN] |

Source and reference attributes

| | |
|---------------------------------|--|
| Submitting organisation: | Australian Commission on Safety and Quality in Health Care |
| Reference documents: | Agency for Clinical Innovation. Key Principles for Care of Confused Hospitalised Older Persons. Sydney: ACI, 2014. Australian Commission on Safety and Quality in Health Care. Delirium Clinical Care Standard. Sydney: ACSQHC; 2021. |