Female—postpartum perineal status, code N[N]

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY4.0 (CC BY4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Female—postpartum perineal status, code N[N]

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Postpartum perineal status
METEOR identifier:	759639
Registration status:	<u>Health</u> , Standard 09/09/2022 <u>Tasmanian Health</u> , Standard 24/03/2023
Definition:	The state of a female's perineum following birth, as represented by a code.
Context:	Perinatal
Data Element Concept:	Female—postpartum perineal status
Value Domain:	Perineal status code N[N]

Value domain attributes

Representational attributes

Representation class:	Code	
Data type:	Number	
Format:	N[N]	
Maximum character length:	2	
	Value	Meaning
Permissible values:	1	Intact
	2	1st degree laceration/vaginal graze
	3	2nd degree laceration
	4	3rd degree laceration
	5	Episiotomy
	7	4th degree laceration
	88	Other perineal laceration, rupture or tear
Supplementary values:	99	Not stated/inadequately described

Collection and usage attributes

Guide for use:

CODE 1 Intact

Should not be recorded in conjunction with any other code.

CODE 2 1st degree laceration/vaginal graze

Graze, laceration, rupture or tear of the perineal skin during birth that may be considered to be slight or that involves one or more of the following structures:

- fourchette
- labia
- periurethral tissue (excluding involvement of urethra)
- vagina, low
- skin
- vulva.

Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

CODE 3 2nd degree laceration

Perineal laceration, rupture or tear as in Code 2 occurring during birth, also involving:

- pelvic floor
- perineal muscles
- vaginal muscles.

Excludes laceration involving the anal sphincter. Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

CODE 4 3rd degree laceration

Perineal laceration, rupture or tear as in Code 3 occurring during birth, also involving:

- anal sphincter (excluding involvement of anal or rectal mucosa)
- rectovaginal septum
- sphincter not otherwise specified (NOS).

Excludes laceration involving the anal or rectal mucosa. Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

CODE 5 Episiotomy

Surgical incision into the perineum and vagina to assist birth.

CODE 7 4th degree laceration

Perineal laceration, rupture or tear as in Code 4 occurring during birth, also involving:

- anal mucosa
- rectal mucosa.

Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

CODE 88 Other perineal laceration, rupture or tear

May include haematoma or unspecified perineal tear. Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

Collection methods:	Jurisdictions that record perinatal data using the ICD-10-AM should apply the following codes in relation to degree of laceration:
	'Code 2 1st degree laceration/vaginal graze' is equivalent to O70.0 in the ICD-10- AM.
	'Code 3 2nd degree laceration ' is equivalent to O70.1 in the ICD-10-AM.
	'Code 4 3rd degree laceration' is equivalent to O70.2 in the ICD-10-AM.
	'Code 7 4th degree laceration' is equivalent to O70.3 in the ICD-10-AM.
	'Code 88 Other perineal laceration, rupture or tear' is equivalent to O70.9 in the ICD-10-AM.
Comments:	Where multiple perinatal lacerations, ruptures or tears of different degrees are documented, assign the code for the highest (most severe) degree only.
	If a laceration occurred during birth and an episiotomy was performed, both the degree of laceration and the episiotomy should be recorded. If an episiotomy is performed, the perineum cannot be intact.
	While 4th degree laceration is more severe than an episiotomy, this category has not been placed in order of clinical significance within the value domain.
	Code 6 (Combined laceration and episiotomy) has been omitted as it is no longer in use. For information about its meaning in previous data elements, see superseded versions.

Source and reference attributes

Submitting organisation:	Australian Institute of Health and Welfare
Origin:	Independent Hospital Pricing Authority 2022. The International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian Modification (ICD-10-AM), Twelfth Edition. Tabular List and Alphabetic Index. Darlinghurst, NSW.

Data element attributes

Collection and usage attributes

Guide for use:	More than one permissible value may be recorded.
Comments:	Perineal laceration (tear) may cause significant maternal morbidity in the postnatal period. Episiotomy is an indicator of management during labour and, to some extent, of obstetric intervention.

Source and reference attributes

Submitting organisation:	Australian Institute of Health and Welfare
--------------------------	--

Relational attributes

Related metadata	Supersedes Female—postpartum perineal status, code N[N]
references:	Health, Superseded 09/09/2022

Specifications:

Implementation in Data Set Mother delivery related data elements (TDLU) cluster Tasmanian Health, Standard 17/11/2023 Implementation start date: 01/07/2023 Implementation end date: 30/06/2025

> Perinatal NMDS 2022-23 Health, Superseded 09/12/2022 Implementation start date: 01/07/2022 Implementation end date: 30/06/2023 DSS specific information:

This is a multiple response data element and is therefore operationalised for data collection across 7 individual data items (one data item per permissible value, excluding Code 99 (Not stated/inadequately described)).

This data element is recorded for the mother only.

Perinatal NMDS 2023-24

Health, Superseded 06/12/2023 Implementation start date: 01/07/2023 Implementation end date: 30/06/2024 DSS specific information:

This is a multiple response data element and is therefore operationalised for data collection across 7 individual data items (one data item per permissible value, excluding Code 99 (Not stated/inadequately described)).

This data element is recorded for the mother only.

Perinatal NMDS 2024–25 Health, Standard 06/12/2023 Implementation start date: 01/07/2024 Implementation end date: 30/06/2025 DSS specific information:

This is a multiple response data element and is therefore operationalised for data collection across 7 individual data items (one data item per permissible value, excluding Code 99 (Not stated/inadequately described)).

This data element is recorded for the mother only.

Tasmanian Perinatal Data Set - 2023 Tasmanian Health, Standard 23/11/2023 Implementation start date: 01/07/2023 Implementation end date: 30/06/2024

Implementation in Indicators:

Used as Numerator

National Core Maternity Indicators: PI03–Episiotomy for females giving birth for the first time and giving birth vaginally, 2024 Health, Recorded 21/02/2024

National Core Maternity Indicators: PI 13-Third and fourth degree tears for vaginal births, 2024

Health, Recorded 21/02/2024

Used as Disaggregation

National Core Maternity Indicators: PI 13-Third and fourth degree tears for vaginal births, 2024

Health, Recorded 21/02/2024