

Australian Health Performance Framework: PI 2.2.2— Healthcare-associated Staphylococcus aureus bloodstream infections, 2022

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Australian Health Performance Framework: PI 2.2.2– Healthcare-associated Staphylococcus aureus bloodstream infections, 2022

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	AHPF PI 2.2.2–Healthcare-associated Staphylococcus aureus bloodstream infections, 2022
Synonymous names:	Healthcare associated Staphylococcus aureus bacteraemia; Healthcare associated Staphylococcus aureus bloodstream infections
METEOR identifier:	778297
Registration status:	Health , Recorded 29/04/2024
Description:	<i>Staphylococcus aureus</i> bloodstream infection (SABSI) associated with care in public hospitals (excluding cases associated with private hospitals and non-hospital care).
Indicator set:	Australian Health Performance Framework, 2022 Health , Recorded 23/04/2024

Collection and usage attributes

Computation description: For the purpose of data collection, all types of public hospitals are included (as defined in the [Local Hospital Networks/Public hospital establishments NMDS 2021–22](#)), both those focusing on acute care, and those focusing on non-acute or sub-acute care, including psychiatric, rehabilitation and palliative care.

Unqualified newborns, hospital boarders and posthumous organ procurement are excluded from the denominator of the indicator.

A patient-episode of SABSIs is defined as a positive blood culture for [Staphylococcus aureus](#). For surveillance purposes, only the first isolate per patient is counted, unless at least 14 days has passed without a positive blood culture, after which an additional episode is recorded.

A *Staphylococcus aureus* bloodstream infection will be considered to be healthcare-associated if: the first positive blood culture is collected more than 48 hours after hospital admission or less than 48 hours after discharge, OR, if the first positive blood culture is collected less than or equal to 48 hours after admission to hospital and the patient-episode of SABSIs meets at least one of the following criteria:

1. SABSIs is a complication of the presence of an indwelling medical device (e.g. intravascular line, haemodialysis vascular access, cerebrospinal fluid (CSF) shunt, urinary catheter)
2. SABSIs occurs within 30 days of a surgical procedure where the SABSIs is related to the surgical site
3. SABSIs was diagnosed within 48 hours of a related invasive instrumentation or incision
4. SABSIs is associated with neutropenia contributed to by cytotoxic therapy. Neutropenia is defined as at least 2 separate calendar days with values of absolute neutrophil count (ANC) or total white blood cell count (WBC) <500 cells/mm³ (0.5×10^9 / L) on or within a 7-day time period which includes the date the positive blood specimen was collected (Day 1), the 3 calendar days before and the 3 calendar days after.

Exclusions:

Cases where a known previous positive test has been obtained within the last 14 days are excluded. For example: if a patient has SABSIs in which 4 sets of blood cultures are positive over the initial 3 days of the patient's admission only one episode of SABSIs is recorded. If the same patient had a further set of positive blood cultures on day 6 of the same admission, these would not be counted again, but would be considered part of the initial patient-episode.

Note: If the same patient had a further positive blood culture 20 days after admission (i.e. greater than 14 days after their last positive blood culture on day 5), then this would be considered a second patient-episode of SABSIs.

See [Establishment—number of patient days, total N\[N\(7\)\]](#) for the definition of patient days.

Analysis by state and territory is based on location of the hospital.

Presented as:

- a number, and
- per 10,000 patient days.

Coverage: Denominator ÷ Number of patient days for all public hospitals in the state or territory.

Any variation from the specifications by jurisdictions will be footnoted and described in the [data quality statement for the National Staphylococcus aureus Bacteraemia Data Collection \(NSABDC\)](#).

Computation:

Numerator

(Numerator ÷ Denominator) x 10,000

Numerator: Number of SABSI patient episodes (as defined above) associated with acute care public hospitals.

Numerator data elements:

Data Element / Data Set

[Establishment—number of healthcare-associated Methicillin-resistant Staphylococcus aureus \(MRSA\) bloodstream infection patient episodes, total episodes N\[NNNN\]](#)

Data Source

[State/territory infection surveillance data](#)

NMDS / DSS

[Healthcare-associated infections NBEDS 2021–](#)

Guide for use

Data source type: Administrative by-product data

Data Element / Data Set

[Establishment—number of healthcare-associated Methicillin-sensitive Staphylococcus aureus \(MSSA\) bloodstream infection patient episodes, total episodes N\[NNNN\]](#)

Data Source

[State/territory infection surveillance data](#)

NMDS / DSS

[Healthcare-associated infections NBEDS 2021–](#)

Guide for use

Data source type: Administrative by-product data

Data Element / Data Set

[Establishment—Staphylococcus aureus bloodstream infection surveillance indicator, yes/no code N](#)

Data Source

[State/territory infection surveillance data](#)

NMDS / DSS

[Healthcare-associated infections NBEDS 2021–](#)

Guide for use

Data source type: Administrative by-product data

Denominator: Number of patient days for acute care public hospitals under surveillance (i.e., only for hospitals included in the surveillance arrangements).

Exclude unqualified newborns, posthumous organ procurement and hospital boarders.

Denominator data elements:

Data Element / Data Set

[Establishment—Staphylococcus aureus bloodstream infection surveillance indicator, yes/no code N](#)

Data Source

[State/territory admitted patient data](#)

NMDS / DSS

[Healthcare-associated infections NBEDS 2021–](#)

Guide for use

Data source type: Administrative by-product data

Data Element / Data Set

[Establishment—number of hospital patient days under Staphylococcus aureus bloodstream infection surveillance, total days N\[NNNNN\]](#)

Data Source

[State/territory admitted patient data](#)

NMDS / DSS

[Healthcare-associated infections NBEDS 2021–](#)

Guide for use

Data source type: Administrative by-product data

Disaggregation:

2017–18 (updated for resupplied data), 2021–22—State and territory, by:

- Methicillin-resistant *Staphylococcus aureus* (MRSA)/Methicillin-sensitive *Staphylococcus aureus* (MSSA)

Some disaggregation may result in numbers too small for publication.

Disaggregation data elements:

Data Element / Data Set

[Establishment—Australian state/territory identifier, code N](#)

Data Source

[State/territory infection surveillance data](#)

NMDS / DSS

[Healthcare-associated infections NBEDS 2021–](#)

Guide for use

Data source type: Administrative by-product data

Comments:

Most recent data available for 2022 Australian Health Performance Framework reporting: 2021–22.

The number of SABS patient episodes associated with acute care public hospitals under surveillance includes SABS patient episodes associated with all public hospitals, and the number of patient days for acute care public hospitals under surveillance includes the number of patient days for all public hospitals under surveillance.

For some states and territories there is less than 100% coverage of hospitals. This may impact on the reported rate. For those jurisdictions with incomplete coverage

of acute care public hospitals (in the numerator), only patient days for those hospitals that contribute data are included (in the denominator). Specifically, if a hospital was not included in the SABSİ surveillance arrangements for part of the year, then the patient days for that part of the year are excluded. If part of the hospital was not included in the SABSİ surveillance arrangements (e.g. children's wards, psychiatric wards), then patient days for that part of the hospital are excluded. Patient days for 'non-acute' hospitals (such as rehabilitation and psychiatric hospitals) are included if the hospital was included in the SABSİ surveillance arrangements, but not otherwise. However, all these patient days are included in the coverage rate denominator measure of total number of patient days for all public hospitals in a state or territory.

Some states operate a 'signal surveillance' arrangement for smaller hospitals whereby the hospital notifies the appropriate authority if a SABSİ case is identified, but the hospital is not considered to have formal SABSİ surveillance as per larger hospitals. Where this arrangement is in place, these hospitals should be included as part of the indicator. That is, SABSİ patient episodes and patient days should be included as 'under surveillance'.

Only episodes associated with acute care public hospital care in each jurisdiction should be counted. If a case is associated with care provided in another jurisdiction (cross border flows) then it is reported (where known) by the jurisdiction where the care associated with the SABSİ occurred.

There may be patient episodes of SABSİ identified by a hospital, which did not originate in the identifying hospital (as determined by the definition of a patient episode of SABSİ), but in another public hospital. If the originating hospital is under SABSİ surveillance, then the patient episode of SABSİ should be attributed to the originating hospital and should be included as part of the indicator. If the originating hospital is not under SABSİ surveillance, then the patient episode is unable to be included in the indicator.

Patient episodes associated with care provided by private hospitals and non-hospital health care are excluded.

Patient days for unqualified newborns, hospital boarders and posthumous organ procurement are excluded.

Almost all patient episodes of SABSİ will be diagnosed when the patient is an admitted patient. However, the intention is that cases are reported whether they were associated with admitted patient care or non-admitted patient care in acute care public hospitals.

Where there is significant variation, for example non-coverage of cases diagnosed less than 48 hours after admission, in the data collection arrangements it will affect the calculation of values across states and territories.

Variation in admission practices across jurisdictions will influence the denominator for this indicator impacting on comparability of rates.

Jurisdictional manuals should be referred to for full details of definitions used in infection control surveillance.

Note that the definition of a healthcare-associated SABSİ was revised by the Australian Commission on Safety and Quality in Health Care in 2016. In particular, the clinical criterion for SABSİ associated with neutropenia was revised. Data from 2010–11 to 2014–15 are reported according to the previous neutropenia criterion:

- SABSİ is associated with neutropenia ($<1 \times 10^9$) contributed to by cytotoxic therapy

Data from 2015–16 onwards are reported according to the new neutropenia criterion:

- SABSİ is associated with neutropenia contributed to by cytotoxic therapy. Neutropenia is defined as at least 2 separate calendar days with values of absolute neutrophil count (ANC) or total white blood cell count (WBC) <500 cell/mm³ (0.5×10^9 /L) on or within a 7-day time period which includes the date the positive blood specimen was collected (Day 1), the 3 calendar days before and the 3 calendar days after.

Note that patient episodes of SABSI are just one type of healthcare associated infection. Hence, this performance indicator is not a complete measure of healthcare associated infections for the outcome area of Hospital and Related Care.

Representational attributes

Representation class: Rate
Data type: Real
Unit of measure: Episode
Format: N[NN].N

Data source attributes

Data sources:

Data Source

[State/territory admitted patient data](#)

Frequency

Annual

Data custodian

State/territory health authorities

Data Source

[State/territory infection surveillance data](#)

Frequency

Annual

Data custodian

State/territory health authorities

Accountability attributes

Reporting requirements: Australian Health Performance Framework

Organisation responsible for providing data: Australian Institute of Health and Welfare

Accountability: Australian Institute of Health and Welfare

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare

Relational attributes

Related metadata references: Supersedes [Australian Health Performance Framework: PI 2.2.2–Healthcare-associated Staphylococcus aureus bloodstream infections, 2021](#)
Health, Standard 11/07/2023

See also [National Healthcare Agreement: PI 22–Healthcare associated infections: Staphylococcus aureus bacteraemia, 2022](#)
Health, Standard 24/09/2021