

Hip Fracture Clinical Care Standard: 2a-Proportion of patients with a hip fracture who either received analgesia within 30 minutes of presentation or did not require it according to an assessment of their pain

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Hip Fracture Clinical Care Standard: 2a-Proportion of patients with a hip fracture who either received analgesia within 30 minutes of presentation or did not require it according to an assessment of their pain

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	2a-Proportion of patients with a hip fracture who either received analgesia within 30 minutes of presentation or did not require it according to an assessment of their pain
METEOR identifier:	780825
Registration status:	Australian Commission on Safety and Quality in Health Care, Standard 10/09/2023
Description:	Proportion of patients with a hip fracture who either received analgesia within 30 minutes of presentation or did not require it according to an assessment of their pain.
Indicator set:	Hip Fracture Clinical Care Standard 2023 Australian Commission on Safety and Quality in Health Care, Standard 10/09/2023

Collection and usage attributes

Population group age from:	50 years or older
Computation description:	To be included in the numerator, patients must have a documented outcome of a pain assessment in their medical record AND either: <ul style="list-style-type: none">• Received analgesia within 30 minutes of presenting to hospital, OR• They did not require analgesia according to their pain assessment. <p>Presented as a percentage.</p>
Computation:	$(\text{Numerator} \div \text{denominator}) \times 100$
Numerator:	Number of patients with a hip fracture who received analgesia within 30 minutes of presentation to hospital or did not require it according to an assessment of their pain.
Denominator:	Number of patients with a hip fracture who presented to hospital.

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Service event
Format:	N[NN]

Accountability attributes

Reporting requirements:	Hip Fracture Clinical Care Standard 2023
Organisation responsible for providing data:	Health service organisations
Accountability:	Health service organisations

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Australian Commission on Safety and Quality in Health Care. Hip Fracture Clinical Care Standard. Sydney: ACSQHC; 2023.

Relational attributes

Related metadata references: Supersedes [Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it](#)
[Health](#), Standard 12/09/2016