Hip Fracture Clinical Care Standard: 2a-Proportion of patients with a hip fracture who either received analgesia within 30 minutes of presentation or did not require it according to an assessment of their pain



© Australian Institute of Health and Welfare 2024

This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Hip Fracture Clinical Care Standard: 2a-Proportion of patients with a hip fracture who either received analgesia within 30 minutes of presentation or did not require it according to an assessment of their pain

Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

Short name: 2a-Proportion of patients with a hip fracture who either received analgesia within

30 minutes of presentation or did not require it according to an assessment of their

pain

METEOR identifier: 780825

Registration status: Australian Commission on Safety and Quality in Health Care, Standard 10/09/2023

Description: Proportion of patients with a hip fracture who either received analgesia within 30

minutes of presentation or did not require it according to an assessment of their

pain.

Indicator set: Hip Fracture Clinical Care Standard 2023

Australian Commission on Safety and Quality in Health Care, Standard

10/09/2023

Collection and usage attributes

Population group age

from:

50 years or older

Computation description: To be included in the numerator, patients must have a documented outcome of a

pain assessment in their medical record AND either:

• Received analgesia within 30 minutes of presenting to hospital, OR

• They did not require analgesia according to their pain assessment.

Presented as a percentage.

Computation: (Numerator ÷ denominator) x 100

Numerator: Number of patients with a hip fracture who received analgesia within 30 minutes of

presentation to hospital or did not require it according to an assessment of their

pain.

Denominator: Number of patients with a hip fracture who presented to hospital.

Representational attributes

Representation class: Percentage

Data type: Real

Unit of measure: Service event

Format: N[NN]

Accountability attributes

Reporting requirements: Hip Fracture Clinical Care Standard 2023

Organisation responsible

for providing data:

Health service organisations

Accountability:

Health service organisations

Metadata 780825 Page 2 of 3 Downloaded 27-Apr-2024

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Australian Commission on Safety and Quality in Health Care. Hip Fracture Clinical

Care Standard. Sydney: ACSQHC; 2023.

Relational attributes

Related metadata references:

Supersedes Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time

or do not require it

Health, Standard 12/09/2016