

Hip Fracture Clinical Care Standard: 3a-Proportion of patients with a hip fracture who had a clinical frailty assessment using a validated tool

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Hip Fracture Clinical Care Standard: 3a-Proportion of patients with a hip fracture who had a clinical frailty assessment using a validated tool

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	3a-Proportion of patients with a hip fracture who had a clinical frailty assessment using a validated tool.
METEOR identifier:	780847
Registration status:	Australian Commission on Safety and Quality in Health Care, Standard 10/09/2023
Description:	Proportion of patients with a hip fracture who had a clinical frailty assessment using a validated tool.
Indicator set:	Hip Fracture Clinical Care Standard 2023 Australian Commission on Safety and Quality in Health Care, Standard 10/09/2023

Collection and usage attributes

Population group age from:	50 years or older
Computation description:	To be included in the numerator, the outcome of the frailty assessment must be document in the medical record. An example of a validated tool is the Clinical Frailty Scale (Rockwood, 2005).
Numerator:	The number of patients with a hip fracture who had a clinical frailty assessment using a validated tool while in hospital.
Denominator:	The number of patients with a hip fracture who were admitted to hospital.

Representational attributes

Representation class:	Percentage
Data type:	Real
Unit of measure:	Person
Format:	N[NN]

Accountability attributes

Reporting requirements:	Hip Fracture Clinical Care Standard 2023
Organisation responsible for providing data:	Health Service Organisations
Accountability:	Health Service Organisations

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
Reference documents:	Australian Commission on Safety and Quality in Health Care. Hip Fracture Clinical Care Standard. Sydney: ACSQHC; 2023. Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, Mitnitski A. A global clinical measure of fitness and frailty in elderly people. CMAJ. 2005 Aug 30;173(5):489-95.

